



Pearson
Edexcel

Mark Scheme (Results)

November 2021

Pearson Edexcel GCE
In Mathematics (9MA0)
Paper 32 Mechanics

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2021

Question Paper Log Number P68824A

Publications Code 9MA0_03_2111_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 100.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response.
If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.


6. Ignore wrong working or incorrect statements following a correct answer.
7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternative answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

General Principles for Mechanics Marking

(But note that specific mark schemes may sometimes override these general principles)

- Rules for M marks: correct no. of terms; dimensionally correct; all terms that need resolving (i.e. multiplied by cos or sin) are resolved.
- Omission or extra g in a resolution is an accuracy error not method error.
- Omission of mass from a resolution is a method error.
- Omission of a length from a moments equation is a method error.
- Omission of units or incorrect units is not (usually) counted as an accuracy error.
- dM indicates a dependent method mark i.e. one that can only be awarded if a previous specified method mark has been awarded.
- Any numerical answer which comes from use of $g = 9.8$ should be given to 2 or 3 SF.
- Use of $g = 9.81$ should be penalised once per (complete) question.
N.B. Over-accuracy or under-accuracy of correct answers should only be penalised *once* per complete question. However, premature approximation should be penalised every time it occurs.
- Marks must be entered in the same order as they appear on the mark scheme.
- In all cases, if the candidate clearly labels their working under a particular part of a question i.e. (a) or (b) or (c),.....then that working can only score marks for that part of the question.
- Accept column vectors in all cases.
- Misreads – if a misread does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, bearing in mind that after a misread, the subsequent A marks affected are treated as A ft
- Mechanics Abbreviations
 - M(A) Taking moments about A
 - N2L Newton's Second Law (Equation of Motion)
 - NEL Newton's Experimental Law (Newton's Law of Impact)
 - HL Hooke's Law
 - SHM Simple harmonic motion
 - PCLM Principle of conservation of linear momentum
 - RHS, LHS Right hand side, left hand side

Question	Scheme		Marks	AOs
1(a)	Use of $\mathbf{v} = \mathbf{u} + \mathbf{at}$ with $t = 2$: $\mathbf{v} = 4\mathbf{i} + 2(2\mathbf{i} - 3\mathbf{j})$		M1	3.1a
	OR integration: $\mathbf{v} = (2\mathbf{i} - 3\mathbf{j})t + 4\mathbf{i}$, with $t = 2$			
	$\mathbf{v} = 8\mathbf{i} - 6\mathbf{j}$		A1	1.1b
			(2)	
1(b)	Use of $\mathbf{r} = \mathbf{ut} + \frac{1}{2}\mathbf{at}^2$ at $t = 3$:			
	$(\mathbf{i} + \mathbf{j}) + \left[3 \times 4\mathbf{i} + \frac{1}{2} \times (2\mathbf{i} - 3\mathbf{j}) \times 3^2 \right]$			
	OR: find \mathbf{v} at $t = 3$: $4\mathbf{i} + 3(2\mathbf{i} - 3\mathbf{j}) = (10\mathbf{i} - 9\mathbf{j})$			
	then use $\mathbf{r} = \frac{1}{2}(\mathbf{u} + \mathbf{v})t$			
	$(\mathbf{i} + \mathbf{j}) + \left[\frac{1}{2} [4\mathbf{i} + (10\mathbf{i} - 9\mathbf{j})] \times 3 \right]$		M1	3.1a
	or $\mathbf{r} = \mathbf{vt} - \frac{1}{2}\mathbf{at}^2$			
	$(\mathbf{i} + \mathbf{j}) + \left[3 \times (10\mathbf{i} - 9\mathbf{j}) - \frac{1}{2} \times (2\mathbf{i} - 3\mathbf{j}) \times 3^2 \right]$			
	OR integration: $\mathbf{r} = (\mathbf{i} + \mathbf{j}) + \left[(2\mathbf{i} - 3\mathbf{j}) \frac{1}{2}t^2 + 4\mathbf{i} \right]$, with $t = 3$			
	$\mathbf{r} = 22\mathbf{i} - 12.5\mathbf{j}$		A1	2.2a
			(2)	
(4 marks)				
Notes: Accept column vectors throughout				
1a	M1	Complete method to find \mathbf{v} , using \mathbf{ruvat} or integration (M0 if \mathbf{i} and/or \mathbf{j} is missing)		
	A1	Apply isw if they also find the speed		
1b	M1	Complete method to find the p.v. but this mark can be scored if they omit $(\mathbf{i} + \mathbf{j})$ i.e. the M1 is for the expression in the square bracket If they integrate, the M1 is earned once the expression in the square bracket is seen with $t = 3$ (M0 if \mathbf{i} and/or \mathbf{j} is missing)		
	A1	cao		

Question	Scheme	Marks	AOs
	Mark parts (a) and (b) together		
2(a)	Equation of motion for A	M1	3.3
	$3mg \sin \alpha - F - T = 3ma$	A1	1.1b
		(2)	
2(b)	Resolve perpendicular to the plane	M1	3.4
	$R = 3mg \cos \alpha$	A1	1.1b
	$F = \frac{1}{6}R$	B1	1.2
	Equation of motion for B OR for whole system	M1	3.3
	$T - mg = ma$ OR $3mg \sin \alpha - F - mg = 3ma + ma$	A1	1.1b
	Complete method to solve for a	DM1	3.1b
	$a = \frac{1}{10}g$ *	A1*	2.2a
		(7)	
2(c)		B1	1.1b
	e.g. acceleration (of B) is constant; dependent on first B1	DB1	2.4
		(2)	
2(d)	e.g. the tensions in the two equations of motion would be different. Tension on A would be different to tension on B	B1	3.5a
		(1)	
(12 marks)			
Notes: N.B. If m's are consistently missing treat as a MR, so max (a) M1A0 (b) M1A0B0M1A1M1A1 (c) B1B1 (d) B1			
For (a) and (b), allow verification, but must see full equations of motion.			
2a	M1	Equation in T and a with correct no. of terms, condone sign errors and sin/cos confusion (If one of the 3's is missing, allow M1) N.B. Treat sin(3/5) etc as an A error but allow recovery	
	A1	Correct equation (allow $(-a)$ instead of a in <u>both</u> equations)	

2b	M1	Correct no. of terms, condone sign errors and sin/cos confusion Allow if appears in (a)
	A1	Correct equation
	B1	Seen anywhere in (a) or (b), including on a diagram
	M1	Equation (for B) in T and a with correct no. of terms, condone sign errors and sin/cos confusion OR Whole system equation with correct no. of terms, condone sign errors and sin/cos confusion
	A1	Correct equation
	DM1	Complete method (trig may not be substituted), dependent on M1 in (a) and second M1 in (b) if they use two equations, or second M1 in (b) if they use one equation.
	A1*	Correct answer correctly obtained.
2c	B1	Straight line starting at the origin (could be reflected in the t -axis). B0 if continuous vertical line at the end.
	DB1	Dependent on first B1, for any equivalent statement
2d	B1	B0 if incorrect extras

Question	Scheme	Marks	AOs
	Part (a) is a 'Show that..' so equations need to be given in full to earn A marks		
3(a)			
	Moments equation: (M1A0 for a moments inequality)	M1	3.3
	$M(A), mga \cos \theta = 2Sa \sin \theta$ $M(B), mga \cos \theta + 2Fa \sin \theta = 2Ra \cos \theta$ $M(C), F \times 2a \sin \theta = mga \cos \theta$ $M(D), 2Ra \cos \theta = mga \cos \theta + 2Sa \sin \theta$ $M(G), Ra \cos \theta = Fa \sin \theta + Sa \sin \theta .$	A1	1.1b
	$(\updownarrow) R = mg$ OR $(\leftrightarrow) F = S$	B1	3.4
	Use their equations (<u>they must have enough</u>) and $F \leq \mu R$ to give an inequality in μ and θ only (allow DM1 for use of $F = \mu R$ to give an equation in μ and θ only)	DM1	2.1
	$\mu \geq \frac{1}{2} \cot \theta^*$	A1*	2.2a
	(5)		
3(b)			
	Moments equation:	M1	3.4
	$M(A), mga \cos \theta = 2Na \sin \theta$ $M(B), mga \cos \theta + 2kmga \sin \theta = 2Ra \cos \theta + \frac{1}{2} mg 2a \sin \theta$ $M(D), 2Ra \cos \theta = mga \cos \theta + N 2a \sin \theta$ $M(G), kmg a \sin \theta + Na \sin \theta = \frac{1}{2} mga \sin \theta + Ra \cos \theta$	A1	1.1b

		$\text{S.C. M}(C), mga \cos \theta + \frac{1}{2}mg2a \sin \theta = kmg2a \sin \theta \quad \mathbf{M1A1B1}$ $1 + \frac{5}{4} = \frac{5k}{2} \quad \mathbf{M1}$ $k = 0.9 \quad \mathbf{A1}$		
		$N = kmg - F \quad \mathbf{OR} \quad R = mg$	B1	3.3
		Use their equations (<u>they must have enough</u>) to solve for k (numerical)	DM1	3.1b
		$k = 0.9 \quad \text{oe}$	A1	1.1b
			(5)	
(10 marks)				
Notes:				
3a	M1	Any moments equation with correct terms, condone sign errors and sin/cos confusion		
	A1	Correct equation		
	B1	Correct equation		
	DM1	Dependent on M1, for using their equations (<u>they must have enough</u>) and $F \leq \mu R$ to give an inequality in μ and θ only (allow M1 for use of $F = \mu R$ to give an equation in μ and θ only)		
	A1*	Given answer correctly obtained with no wrong working seen (e.g. if they use $F = \mu R$ anywhere, A0)		
3b	M1	Any moments equation with correct terms, condone sign errors		
	A1	Correct equation		
	B1	Correct equation		
	DM1	Dependent on M1, for using their equations (<u>they must have enough</u>) with trig substituted, to solve for k , which must be numerical.		
	A1	cao		

Question	Scheme		Marks	AOs
	Note that $g = 10$; penalise once for whole question if $g = 9.8$			
4(a)	Use $s = ut + \frac{1}{2}at^2$ vertically or any complete method to give an equation in t only		M1	3.4
	$-70 = 65 \sin \alpha \times t - \frac{1}{2} \times g \times t^2$		A1	1.1b
			M(A)1	1.1b
	$t = 7$ (s)		A1	1.1b
			(4)	
4(b)	Horizontal velocity component at $A = 65 \cos \alpha$ (60)		B1	3.4
	Complete method to find vertical velocity component at A		M1	3.4
	$65 \sin \alpha - g \times 7$ OR $\sqrt{(-25)^2 + 2g \times 70}$ (45)		A1ft	1.1b
	Sub for trig and square, add and square root : $\sqrt{60^2 + (-45)^2}$		M1	3.1b
	75 Accept 80 (m s^{-1})		A1	1.1b
				(5)
4(c)	e.g. an approximate value of g has been used, the dimensions of the stone could affect its motion, spin of the stone, $g = 10$ instead of 9.8 has been used, g has been assumed to be constant, wind effect, shape of the stone		B1	3.5b
			(1)	
(10 marks)				
Notes:				
4a	M1	Complete method, correct no. of terms, condone sign errors and sin/cos confusion		
	A1	Correct equation in t only with at most one error		
	M(A)1	Correct equation in t only		
		N.B. For ‘up and down’ methods etc, the two A marks are for all the equations that they use, lose a mark for each error.		
	A1	Cao ($g = 9.8, 7.1$ or 7.11) ($g = 9.81, 7.1$ or 7.12)		
4b	B1	Seen, including on a diagram.		
	M1	Condone sign errors and sin/cos confusion		
	A1ft	Correct expression; accept negative of this, follow their t		
	M1	Sub for trig and use Pythagoras		
	A1	Cao ($g = 9.8$ or $9.81, 75$ or 74.8)		

4c	B1	B0 if incorrect extras
----	----	------------------------

Question	Scheme	Marks	AOs
	Allow column vectors throughout this question		
5(a)	Differentiate \mathbf{v} wrt t	M1	3.1a
	$\frac{3}{2}t^{-\frac{1}{2}}\mathbf{i} - 2\mathbf{j}$ isw	A1	1.1b
		(2)	
5(b)	$3t^{\frac{1}{2}} = 2t$	M1	2.1
	Solve for t	DM1	1.1b
	$t = \frac{9}{4}$	A1	1.1b
		(3)	
5(c)	Integrate \mathbf{v} wrt t	M1	3.1a
	$\mathbf{r} = 2t^{\frac{3}{2}}\mathbf{i} - t^2\mathbf{j} (+\mathbf{C})$	A1	1.1b
	$t = 1, \mathbf{r} = -\mathbf{j} \Rightarrow \mathbf{C} = -2\mathbf{i}$ so $\mathbf{r} = 2t^{\frac{3}{2}}\mathbf{i} - t^2\mathbf{j} - 2\mathbf{i}$	A1	2.2a
		(3)	
5(d)	$\sqrt{(3t^{\frac{1}{2}})^2 + (2t)^2} = 10$ or $(3t^{\frac{1}{2}})^2 + (2t)^2 = 10^2$	M1	2.1
	$9t + 4t^2 = 100$	M(A)1	1.1b
	$t = 4$	A1	1.1b
	$\mathbf{r} = 14\mathbf{i} - 16\mathbf{j}$	M1	1.1b
	$\sqrt{14^2 + (-16)^2}$	M1	3.1a
	$\sqrt{452} (2\sqrt{113})$ (m)	A1	1.1b
		(6)	

(14 marks)

Notes:

5a	M1	Both powers decreasing by 1 (M0 if vector(s) disappear but allow recovery)
	A1	cao
5b	M1	Complete method, using \mathbf{v} , to obtain an equation in t only, allow a sign error
	DM1	Dependent on M1, solve for t

	A1	cao
5c	M1	Both powers increasing by 1 (M0 if vectors disappear but allow recovery)
	A1	Correct expression without C
	A1	cao
5d	M1	Use of Pythagoras on v and 10 to set up equation in <i>t</i>
	M(A)1	Correct 3 term quadratic in <i>t</i>
	A1	cao
	M1	Substitute their numerical <i>t</i> value into their r
	M1	Use of Pythagoras to find the magnitude of their r
	A1	cso